
PGC7, H3K9me2 and Tet3: regulators of DNA methylation in zygotes.

Journal: Cell Res

Publication Year: 2013

Authors: Jinsuk Kang, Sundeep Kalantry, Anjana Rao

PubMed link: 22868271

Funding Grants: Generation of regulatory T cells by reprogramming

Public Summary:

In zygotes, a global loss of DNA methylation occurs selectively in the paternal pronucleus before the first cell division, concomitantly with the appearance of modified forms of 5-methylcytosine. The adjacent maternal pronucleus and certain paternally-imprinted loci are protected from this process. Nakamura et al. recently clarified the molecular mechanism involved: PGC7/Stella/Dppa3 binds to dimethylated histone 3 lysine 9 (H3K9me2), thereby blocking the activity of the Tet3 methylcytosine oxidase in the maternal genome as well as at certain imprinted loci in the paternal genome.

Scientific Abstract:

In zygotes, a global loss of DNA methylation occurs selectively in the paternal pronucleus before the first cell division, concomitantly with the appearance of modified forms of 5-methylcytosine. The adjacent maternal pronucleus and certain paternally-imprinted loci are protected from this process. Nakamura et al. recently clarified the molecular mechanism involved: PGC7/Stella/Dppa3 binds to dimethylated histone 3 lysine 9 (H3K9me2), thereby blocking the activity of the Tet3 methylcytosine oxidase in the maternal genome as well as at certain imprinted loci in the paternal genome.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/pgc7-h3k9me2-and-tet3-regulators-dna-methylation-zygotes>